

BRANCHED TWIST SPINS AND KNOT DETERMINANTS

MIZUKI FUKUDA

ABSTRACT. A branched twist spin is a generalization of twist spun knots, which appeared in the study of locally smooth circle actions on the 4-sphere due to Montgomery, Yang, Fintushel and Pao. In this paper, we give a sufficient condition to distinguish non-equivalent, non-trivial branched twist spins by using knot determinants. To prove the assertion, we give a presentation of the fundamental group of the complement of a branched twist spin, which generalizes a presentation of Plotnick, calculate the first elementary ideals and obtain the condition of the knot determinants by substituting -1 for the indeterminate.

1. INTRODUCTION

An n -knot is an n -sphere embedded into the $(n + 2)$ -sphere. In this paper, we assume that all knots are smooth. A non-trivial example of a 2-knot was first introduced by Artin [1], which was constructed from a 1-knot in S^3 by rotating it in S^4 along a trivial axis. It is called a *spun knot*. One of the important properties of spun knots is that its knot group is isomorphic to the knot group of the 1-knot of rotation. Afterwards, Zeeman generalized Artin's construction by adding a twist during the rotation, called a *twist spun knot* [14]. Such a twisting is realized by an S^1 -action on S^4 having two fixed points and one family of exceptional orbits. A twist spun knot is then obtained as the union of the fixed points and the preimage of a (possibly knotted) arc in the orbit space connecting the image of the two fixed points.

Suppose that S^4 has an effective locally smooth S^1 -action. Let E_m be the set of exceptional orbits of \mathbb{Z}_m -type, where m is an integer greater than 1, and F be the fixed point set. Let E_m^* and F^* denote the image of the orbit map of E_m and F , respectively. In the study of S^1 -action on S^4 , Montgomery and Yang [8] showed that effective locally smooth S^1 -actions are classified into the following four types: (1) $\{D^3\}$, (2) $\{S^3\}$, (3) $\{S^3, m\}$, and (4) $\{(S^3, K), m, n\}$, which are called *orbit data*. The 3-ball D^3 and the 3-sphere S^3 in these notations represent the orbit spaces. Type (1) has no exceptional orbit and F^* is the whole boundary of D^3 . Type (2) has no exceptional orbit and F^* consists of two points. Type (3) has one type of exceptional orbits E_m whose image E_m^* constitutes an arc in the orbit space S^3 . The image F^* is the endpoints of the arc E_m^* . Type (4) has two different types of exceptional orbits E_m and E_n , where m, n are larger than 1 and relatively prime. The images E_m^* and E_n^* are arcs with the endpoints F^* such that $E_m^* \cup E_n^* \cup F^*$ constitutes a 1-knot in S^3 . Conversely, Fintushel [5] and Pao [9] showed that there is a unique weak equivalence class of effective locally smooth S^1 -actions on S^4 for each orbit data. Here the weak equivalence of S^1 -manifolds M_1 and M_2 is defined by

2010 *Mathematics Subject Classification*. Primary 57Q45; Secondary 57M60, 57M27.

Key words and phrases. 2-knots, circle actions, Alexander polynomials.

a homeomorphism $H : M_1 \rightarrow M_2$ satisfying $H(\theta x) = a(\theta)H(x)$ for $\theta \in S^1$ and $x \in M_1$, where a is an automorphism of S^1 .

The sets $E_m \cup F$ and $E_n \cup F$ in type (4) are diffeomorphic to 2-spheres and the image of all exceptional orbits and the fixed point set is a 1-knot $K = E_m^* \cup E_n^* \cup F^*$ in S^3 . An (m, n) -branched twist spin of K is defined to be the 2-knot $E_n \cup F$ in the case of (4), which corresponds to the case $m, n > 1$, and defined similarly in type (2) and (3). See Section 1 for precise definition. In this paper, by fixing the orientation of S^4 , we generalize the definition of (m, n) -branched twist spins for $(m, n) \in \mathbb{Z} \times \mathbb{N}$, where $|m|$ and n are coprime. Note that if $n = 1$ then it is of type (3) and corresponds to the m -twist spun knot, and if $m = 0$ and $n = 1$ then it is of type (2) and corresponds to the spun knot.

An n -knot K is said to be equivalent to another n -knot K' , denoted by $K \sim K'$, if there exists a smooth isotopy $H_t : S^{n+2} \rightarrow S^{n+2}$ such that $H_0 = \text{id}$ and $H_1(K) = K'$.

It is known by Hillman and Plotnick that if an 1-knot K is torus or hyperbolic then $K^{m,n}$ is not reflexive for $m \geq n$ and $m \geq 3$ [7]. Here a 2-knot is said to be reflexive if its Gluck reconstruction is equivalent to either its mirror image or its orientation reverse. Since the trivial 2-knot is reflexive, $K^{m,n}$ is non-trivial if K is torus or hyperbolic and $m \geq n$ and $m \geq 3$. Though their result detects non-triviality of these branched twist spins, in the best of my knowledge, there is no result which distinguishes non-equivalent, non-trivial branched twist spins.

In this paper we give a sufficient condition to distinguish non-trivial branched twist spins by using knot determinants. Our main theorem is the following:

Theorem 1.1. Let $K_1^{m_1, n_1}, K_2^{m_2, n_2}$ be branched twist spins constructed from 1-knots K_1 and K_2 in S^3 , respectively.

- (1) If m_1, m_2 are even and $|\Delta_{K_1}(-1)| \neq |\Delta_{K_2}(-1)|$ then $K_1^{m_1, n_1} \not\sim K_2^{m_2, n_2}$.
- (2) If m_1 is even, m_2 is odd and $|\Delta_{K_1}(-1)| \neq 1$ then $K_1^{m_1, n_1} \not\sim K_2^{m_2, n_2}$.

To prove this theorem, we first give a presentation of the fundamental group of the complement of a branched twist spin, which generalizes a presentation of Plotnick, and obtain the first elementary ideal from this presentation by using Fox calculus. Then we obtain certain equations of elementary ideals and the knot determinants appear when we substitute -1 for the variable t of the Alexander polynomials.

This paper is organized as follows: In Section 2, we give the definition of a branched twist spin and the presentation of its fundamental group. In Section 3, we give concrete generators of the first elementary ideal and prove Theorem 1.1.

Acknowledgments. The author is grateful to his advisor, Masaharu Ishikawa, for many helpful suggestions.

2. BRANCHED TWIST SPINS

For a point $x \in M$, an orbit $G(x)$ is called G/H -type if the isotropy group $G_x = \{gx = x \mid g \in G\}$ of x is $H \subset G$. If the G -action is locally smooth, each orbit of G/H -type has a linear slice S such that H acts orthogonally on S . See [3] for the terminologies of transformation groups.

Suppose that S^4 has an effective locally smooth S^1 -action. We consider the S^1 -actions of type (3) and type (4) in Section 1. Let E_m and E_n be the sets of exceptional orbits of

\mathbb{Z}_m -type and \mathbb{Z}_n -type, respectively, and F be the fixed point set. Here, when $n = 1$, the S^1 -action is of type (3). Let E_m^* , E_n^* and F^* denote the image of the orbit map of E_m , E_n and F , respectively. Then $E_m \cup F$ and $E_n \cup F$ are diffeomorphic to 2-spheres and the image of all exceptional orbits and the fixed point set constitutes an 1-knot $K = E_m^* \cup E_n^* \cup F^*$.

Now we give the definition of (m, n) -branched twist spins for $(m, n) \in \mathbb{Z} \times \mathbb{N}$. To do this, we need to fix orientations of S^4 and the S^1 -action on S^4 , and observe the direction of twisting in neighborhoods of the exceptional orbits. Let (m, n) be a pair of integers in $\mathbb{Z} \times \mathbb{N}$ such that $|m|$ and n are coprime. Here we further assume that $m \neq 0$. First, we decompose the orbit space S^3 into five pieces. The set F^* consists of two points, say x_1^* and x_2^* , and let D_i^{3*} be a small compact ball in S^3 centered at x_i^* for $i = 1, 2$. Choose a compact tubular neighborhood $N(K)$ of K sufficiently small such that $N(K) \setminus \text{int}(D_1^{3*} \cup D_2^{3*})$ has two connected components N_m and N_n with $N_m \cap E_m^* \neq \emptyset$ and $N_n \cap E_n^* \neq \emptyset$. Set E_m^{c*} and E_n^{c*} to be the connected components of $N(K) \setminus \text{int}(D_1^{3*} \cup D_2^{3*})$, where $E_m^{c*} \subset N_m$ and $E_n^{c*} \subset N_n$, respectively. Note that N_m and N_n are diffeomorphic to $E_m^{c*} \times D^2$ and $E_n^{c*} \times D^2$, respectively. Let X be the closure of $S^3 \setminus ((E_m^{c*} \times D^2) \cup (E_n^{c*} \times D^2) \cup D_1^{3*} \cup D_2^{3*})$, which is the knot complement of K . Then we have a decomposition $S^3 = X \cup (E_m^{c*} \times D^2) \cup (E_n^{c*} \times D^2) \cup D_1^{3*} \cup D_2^{3*}$.




FIGURE 1. The complement X

Let $p : S^4 \rightarrow S^3$ be the orbit map. Each point of X is the image of a free orbit. Thus $p|X \times S^1$ is a principal S^1 -bundle. The preimage $p^{-1}(X)$ is diffeomorphic to $X \times S^1$ and $p|X \times S^1 : X \times S^1 \rightarrow X$ is the first projection since $H^2(X; \mathbb{Z}) = H_1(X, \partial X; \mathbb{Z}) = 0$ (cf. [2, Chapter 2]).

Let B_i^4 be a linear slice at $p^{-1}(x_i^*)$, which is a closed 4-ball. By [5], $p^{-1}(D_i^{3*})$ is diffeomorphic to B_i^4 and the action on $p^{-1}(D_i^{3*})$ is S^1 -equivalent to that on B_i^4 . Note that the action on B_i^4 is the cone of the action of ∂B_i^4 and so the action on $p^{-1}(D_i^{3*})$ is the cone of the action of $p^{-1}(\partial D_i^{3*})$.

Choosing a point z_m^* in E_m^{c*} , let $D_{z_m^*}^{2*}$ be a 2-disk in S^3 centered at $z_m^* \in E_m^{c*}$ and transversal to E_m^{c*} . The preimage $p^{-1}(D_{z_m^*}^{2*})$ is a solid torus V_m whose core is the exceptional orbit of \mathbb{Z}_m -type. In the same way, choosing a point $z_n^* \in E_n^{c*}$, let $D_{z_n^*}^{2*}$ be a 2-disk in S^3 centered at $z_n^* \in E_n^{c*}$ and transversal to E_n^{c*} . The preimage $p^{-1}(D_{z_n^*}^{2*})$ is a solid torus V_n whose core is the exceptional orbit of \mathbb{Z}_n -type. Note that, since $V_m \cup V_n = p^{-1}(\partial D_i^{3*})$ is a 3-sphere, $p^{-1}(y^*)$ ($y^* \in \partial D_{z_m^*}^{2*}$) is a curve on ∂V_m rotating, up to orientation, m times along the meridian and n times along the longitude of V_m , where n is determined module m due to the self-homeomorphisms of V_m .

Now we fix the orientations of V_m and E_m^{c*} as follows: First, fix the orientation of S^4 and those of orbits such that they coincide with the direction of the S^1 -action. These orientations determine the orientation of $V_m \times E_m^{c*}$. Let (ϕ, θ) be a preferred meridian-longitude pair of X . From the decomposition of the orbit space S^3 , we can see that ϕ is regarded as a coordinate of the second factor of $V_m \times E_m^{c*}$. We assign the orientation of V_m so that the orientation of $V_m \times E_m^{c*}$ coincides with the given orientation of S^4 . Finally, we choose the meridian and longitude pair (Θ, H) of $V_m \cong D^2 \times S^1$ such that H becomes the meridian of V_n in the decomposition $V_m \cup V_n = p^{-1}(\partial D_i^{3*})$ and the orbits of the S^1 -action are in the direction $\varepsilon n\Theta + |m|H$ with $n > 0$, where $\varepsilon = 1$ if $m \geq 0$ and $\varepsilon = -1$ if $m < 0$.




FIGURE 2. The preimage of y^*

Definition 2.1 (Branched twist spin). For each pair $(m, n) \in \mathbb{Z} \times \mathbb{N}$ with $m \neq 0$ such that $|m|$ and n are coprime, let $K^{m,n}$ be the 2-knot $E_n \cup F$. If $(m, n) = (0, 1)$ then let $K^{0,1}$ be the spun knot of K . The 2-knot $K^{m,n}$ is called an (m, n) -branched twist spin of K .

Note that the branched twist spin $K^{m,1}$ constructed from $\{(S^3, K), m, 1\}$ is an m -twist spun knot of K .




FIGURE 3. The image of $E_m^* \cup E_n^* \cup F^*$ in S^3

Let K be an n -knot in S^{n+2} . The fundamental group of the knot complement $S^{n+2} \setminus \text{int}N(K)$ of an n -knot K is called the *knot group* of K , where $N(K)$ is a tubular neighborhood of K .

Lemma 2.2. Let K be an 1-knot and $K^{m,n}$ be the (m, n) -branched twist spin of K with $(m, n) \in \mathbb{Z} \times \mathbb{N}$, where $|m|$ and n are coprime. Let $\langle x_1, \dots, x_s \mid r_1, \dots, r_t \rangle$ be a presentation of the knot group of K such that x_1 is a meridian. Then the knot group of $K^{m,n}$ has the presentation

$$(2.1) \quad \pi_1(S^4 \setminus \text{int}N(K^{m,n})) \cong \langle x_1, \dots, x_s, h \mid r_1, \dots, r_t, x_i h x_i^{-1} h^{-1}, x_1^{|m|} h^\beta \rangle,$$

where β is an integer such that $n\beta \equiv \varepsilon \pmod{m}$. Recall that $\varepsilon = 1$ if $m \geq 0$ and $\varepsilon = -1$ if $m < 0$.

Remark 2.3. The presentation of roll spun knots is similar to the presentation in Lemma 2.2. For twist-roll spun knots, the conditions of distinguishing two twist roll-spins and non-triviality is studied by Teragaito [12, 13].

Proof. The knot complement of $K^{m,n}$ is given by $X \times S^1 \cup_f V_m \times E_m^{c*}$, where the map $f : \partial D^{2*} \times E_m^{c*} \times S^1 \rightarrow \partial V_m \times E_m^{c*}$ is the attaching map specified by the decomposition explained in this section.

Let h is the coordinate of the second factor of $X \times S^1$ whose direction coincides with the S^1 -action. By the above discussion of the orientations, the induced map $f_* : H_1(\partial D^{2*} \times E_m^{c*} \times S^1) \rightarrow H_1(V_m \times E_m^{c*})$ must satisfy

$$(f_*([\theta]), f_*([h])) = ([\Theta], [H]) \begin{pmatrix} \alpha & \varepsilon n \\ -\beta & |m| \end{pmatrix},$$

where α and β are integers satisfying $m\alpha + n\beta = \varepsilon$.

The inverse of f_* satisfies the relation

$$(f_*^{-1}([\Theta]), f_*^{-1}([H])) = ([\theta], [h]) \begin{pmatrix} |m| & -\varepsilon n \\ \beta & \alpha \end{pmatrix},$$

hence $f_*^{-1}([\Theta]) = |m|[\theta] + \beta[h]$ and $f_*^{-1}([H]) = -\varepsilon n[\theta] + \alpha[h]$ hold. Since $f_*^{-1}([\Theta])$ is null-homologous in $V_m \times E_m^{c*}$,

$$\pi_1(S^4 \setminus \text{int}N(K^{m,n})) \cong \langle x_1, \dots, x_s, h \mid r_1, \dots, r_t, x_i h x_i^{-1} h^{-1}, x_1^{|m|} h^\beta \rangle$$

holds by Van Kampen's theorem. \square

Remark 2.4. Since ∂D_i^4 is the union of V_m and V_n , the meridian μ of $K^{m,n}$ is that of V_n , named H in the above proof. From the relation $f_*^{-1}([H]) = -\varepsilon n[\theta] + \alpha[h]$, we have $\mu = \theta^{-\varepsilon n} h^\alpha$.

3. PROOF OF MAIN THEOREM

Let K be an n -knot. Assume that a presentation $\langle x_1, \dots, x_l \mid r_1, \dots, r_k \rangle$ of $\pi_1(S^{n+2} \setminus \text{int}N(K))$ is given. Let $a : \pi_1(X) \rightarrow H_1(X) \cong \mathbb{Z}\langle t \rangle$ be the quotient map. Here $\langle t \rangle$ is the infinite cyclic group. This map induces a map $a_* : \mathbb{Z}\pi_1(X) \rightarrow \mathbb{Z}[t, t^{-1}]$ naturally. The matrix A defined by

$$A = \left(a_* \left(\frac{\partial r_i}{\partial x_j} \right) \right)$$

is called the *Alexander matrix* of K . Note that $H_1(X) \cong \mathbb{Z}$ holds for all n -knots, and a takes a meridian of K to the generator t of \mathbb{Z} .

In the case of a 1-knot, it is very common to use a Wirtinger presentation for describing the knot group of K . Then, the quotient map of the abelianization maps each generator to the generator t of $H_1(X; \mathbb{Z})$.

Two Alexander matrices A and A' is said to be equivalent, denoted by $A \sim A'$, if A' is obtained from A by the following operations: (1)Permuting rows or permuting

columns. (2) Adjoining to a row or a column a linear combination of other rows or columns, respectively. (3) $A \rightarrow \begin{pmatrix} A & \\ & 0 \end{pmatrix}$. (4) $A \rightarrow \begin{pmatrix} A & 0 \\ 0 & 1 \end{pmatrix}$.

For the Alexander matrix $A \in M(p, q, \mathbb{Z}[t, t^{-1}])$ of K over $\mathbb{Z}[t, t^{-1}]$ and non-negative integer k , the k -th elementary ideal $E_k(A)$ of K is defined as follows:

- $E_k(A)$ is the ideal generated by determinants of all $(q - k) \times (q - k)$ -submatrices of A if $0 < q - k \leq p$,
- $E_k(A) = 0$ if $q - k > p$,
- $E_k(A) = \mathbb{Z}[t, t^{-1}]$ if $q - k \leq 0$.

For all k , we have $E_k \subset E_{k+1}$. If $K \sim K'$, a presentation of the knot group of K' is obtained from that of K by Tietze transformations. Since Tietze transformations preserve the equivalence class of Alexander matrices, we denote $E_k(A)$ by $E_k(K)$.

Now we study the elementary ideals of the knot group of $K^{m,n}$. Hereafter we fix a Wirtinger presentation of the knot group K as

$$(3.1) \quad \langle x_1, \dots, x_l \mid r_1, \dots, r_l \rangle.$$

Then,

$$(3.2) \quad \pi_1(S^4 \setminus \text{int}N(K^{m,n})) \cong \langle x_1, \dots, x_l, h \mid r_1, \dots, r_l, x_i h x_i^{-1} h^{-1}, x_1^{|m|} h^\beta \rangle$$

holds by applying Lemma 2.2. Let r_{l+i} be $x_i h x_i^{-1} h^{-1}$ for each $i (1 \leq i \leq l)$, and let r_{2l+1} be $x_1^{|m|} h^\beta$. Then, using the induced map a_* and Fox calculus for this presentation, the Alexander matrix A of $K^{m,n}$ is written as

$$A = \left(a_* \left(\frac{\partial r_i}{\partial x_j} \right) \right).$$

Lemma 3.1. The k -th elementary ideal of $K^{m,n}$ has the following property:

- (1) $E_0(K^{m,n}) = 0$.
- (2) Let β be a positive integer satisfying $n\beta \equiv \varepsilon \pmod{m}$. The ideal $E_1(K^{m,n})$ is the ideal generated by the following elements:

$$\left\{ \begin{array}{l} \Delta_K(t^\beta) \left\{ (1 - t^{|m|}), (1 - t^\beta), \frac{1 - t^{|m|\beta}}{1 - t^\beta}, \frac{1 - t^{|m|\beta}}{1 - t^{|m|}} \right\}, \\ G_i(t^\beta)(1 - t^{|m|}) \left\{ (1 - t^{|m|}), (1 - t^\beta), \frac{1 - t^{|m|\beta}}{1 - t^\beta}, \frac{1 - t^{|m|\beta}}{1 - t^{|m|}} \right\}, \\ (1 - t^{|m|})^{l-1} \left\{ (1 - t^{|m|}), (1 - t^\beta), \frac{1 - t^{|m|\beta}}{1 - t^\beta}, \frac{1 - t^{|m|\beta}}{1 - t^{|m|}} \right\}, \end{array} \right.$$

where l is the number of generators of the knot group of K and $G_i(t)$ are generators of $E_2(K)$. Especially, $E_1(K^{m,n}) \neq 0$.

Here the notation $P\{Q_1, Q_2, Q_3, Q_4\}$ means PQ_1, PQ_2, PQ_3, PQ_4 .

Proof. From Lemma 2.2, we have the presentation (3.2) of the knot group of $K^{m,n}$. By Remark 2.4, the meridian of $K^{m,n}$ is written as $x_1^{-\varepsilon n} h^\alpha$, where $m\alpha + n\beta = \varepsilon$. Since the quotient map a sends $x_1^{-\varepsilon n} h^\alpha$ to the generator of $H_1(X; \mathbb{Z})$ identified with t , we have

$a(x_1) = \cdots = a(x_l) = t^{-\beta}$, $a(h) = t^{|m|}$. Then the Alexander matrix A obtained from (3.2) is given by

$$(3.3) \quad A = \begin{pmatrix} & & & & 0 \\ & & & & \vdots \\ & & & & \vdots \\ & & & & \vdots \\ & & & & 0 \\ & & & & t^{-\beta} - 1 \\ & & & & \vdots \\ & & & & \vdots \\ & & & & \vdots \\ & & & & 0 \\ & & & & t^{-\beta} - 1 \\ & & & & \vdots \\ & & & & \vdots \\ & & & & \vdots \\ & & & & 0 \\ & & & & t^{-|m|\beta}(1-t^{|m|\beta}) \\ & & & & \frac{1-t^{|m|\beta}}{1-t^{|m|}} \end{pmatrix},$$

where B is the Alexander matrix of K obtained from the Wirtinger presentation (3.1) by replaced t with $t^{-\beta}$. Since r_i is $x_i x_j x_i^{-1} x_k^{-1}$, for each i ($1 \leq i \leq l$), entries in the i -th row of A satisfy

$$a_* \left(\frac{\partial x_i x_j x_i^{-1} x_k^{-1}}{\partial x_p} \right) = \begin{cases} 1 - t^{-\beta} & (p = i) \\ t^{-\beta} & (p = j) \\ -1 & (p = k) \\ 0 & (p : \text{others}) \end{cases}.$$

Therefore we have $\sum_{p=1}^l a_* \left(\frac{\partial x_i x_j x_i^{-1} x_k^{-1}}{\partial x_p} \right) = 0$. Hence, A is equivalent to

$$A' = \begin{pmatrix} 0 & & & & 0 \\ \vdots & & & & \vdots \\ \vdots & & & & \vdots \\ \vdots & & & & \vdots \\ 0 & & & & 0 \\ 1 - t^{|m|} & & & & t^{-\beta} - 1 \\ & & & & 0 \\ & & & & \vdots \\ & & & & \vdots \\ & & & & \vdots \\ & & & & 0 \\ & & & & t^{-|m|\beta}(1-t^{|m|\beta}) \\ \frac{1-t^{|m|\beta}}{1-t^{-\beta}} & 0 & \cdots & 0 & \frac{1-t^{|m|\beta}}{1-t^{|m|}} \end{pmatrix},$$

where B_1 is an $l \times (l-1)$ matrix. An $(l+1) \times (l+1)$ submatrix of A' should contain both the $(l+1)$ -th row and the $(2l+1)$ -th row if its determinant is not zero. Then any $(l+1) \times (l+1)$ submatrix has the form

$$\begin{pmatrix} 0 & & & & 0 \\ \vdots & & & & \vdots \\ 0 & & & & 0 \\ 1 - t^{|m|} & 0 & \cdots & 0 & t^{-\beta} - 1 \\ 0 & & & & 0 \\ \vdots & & & & \vdots \\ 0 & & & & 0 \\ \frac{1-t^{|m|\beta}}{1-t^{-\beta}} & 0 & \cdots & 0 & \frac{1-t^{|m|\beta}}{1-t^{|m|}} \end{pmatrix},$$

where $\begin{pmatrix} B_2 \\ B_3 \end{pmatrix}$ is an $(l-1) \times (l-1)$ matrix, and its determinant has the factor

$$\begin{vmatrix} 1 - t^{|m|} & t^{-\beta} - 1 \\ \frac{1 - t^{-|m|\beta}}{1 - t^{-\beta}} & \frac{t^{-|m|\beta}(1 - t^{|m|\beta})}{1 - t^{|m|}} \end{vmatrix}$$

which is equal to zero. Thus $E_0(K^{m,n}) = 0$ holds.

By checking all the determinants of $l \times l$ submatrices of A' , we can see that $E_1(K^{m,n})$ is generated by the following terms:

$$\begin{cases} \Delta_K(t^{-\beta}) \left\{ (1 - t^{|m|}), (t^{-\beta} - 1), \frac{1 - t^{-|m|\beta}}{1 - t^{-\beta}}, \frac{t^{-|m|\beta}(1 - t^{|m|\beta})}{1 - t^{|m|}} \right\}, \\ G_i(t^{-\beta})(1 - t^{|m|}) \left\{ (1 - t^{|m|}), (t^{-\beta} - 1), \frac{1 - t^{-|m|\beta}}{1 - t^{-\beta}}, \frac{t^{-|m|\beta}(1 - t^{|m|\beta})}{1 - t^{|m|}} \right\}, \\ (1 - t^{|m|})^{l-1} \left\{ (1 - t^{|m|}), (t^{-\beta} - 1), \frac{1 - t^{-|m|\beta}}{1 - t^{-\beta}}, \frac{t^{-|m|\beta}(1 - t^{|m|\beta})}{1 - t^{|m|}} \right\}. \end{cases}$$

Here $\Delta_K(t)$ is the Alexander polynomial of K , which is given, up to unit, as the common factor of all determinants of $(l-1) \times (l-1)$ submatrices of B_1 , and $G_i(t)$ are the generators of $E_2(K)$. Thus we have the assertion. \square

Proof of Theorem 1.1. We prove the assertion by contraposition. Suppose that $K_1^{m_1, n_1} \sim K_2^{m_2, n_2}$. Let $G_j^i(t)$ be the generators of $E_2(K_i)$. From Lemma 3.1, for each $i = 1, 2$, $E_1(K^{m_i, n_i})$ is generated by

$$\begin{cases} \Delta_{K_i}(t^{\beta_i}) \left\{ (1 - t^{|m_i|}), (1 - t^{\beta_i}), \frac{1 - t^{|m_i|\beta_i}}{1 - t^{\beta_i}}, \frac{1 - t^{|m_i|\beta_i}}{1 - t^{|m_i|}} \right\}, \\ G_j^i(t^{\beta_i})(1 - t^{|m_i|}) \left\{ (1 - t^{|m_i|}), (1 - t^{\beta_i}), \frac{1 - t^{|m_i|\beta_i}}{1 - t^{\beta_i}}, \frac{1 - t^{|m_i|\beta_i}}{1 - t^{|m_i|}} \right\}, \\ (1 - t^{|m_i|})^{l_i-1} \left\{ (1 - t^{|m_i|}), (1 - t^{\beta_i}), \frac{1 - t^{|m_i|\beta_i}}{1 - t^{\beta_i}}, \frac{1 - t^{|m_i|\beta_i}}{1 - t^{|m_i|}} \right\}. \end{cases}$$

Since the ideals $E_1(K_1^{m_1, n_1})$ and $E_1(K_2^{m_2, n_2})$ coincide, each generator of $E_1(K_1^{m_1, n_1})$ is a linear combination of generators of $E_1(K_2^{m_2, n_2})$ over $\mathbb{Z}[t, t^{-1}]$. Thus, for instance, we have

$$\begin{aligned} \Delta_{K_2}(t^{\beta_2})(1 - t^{|m_2|}) &= \Delta_{K_1}(t^{\beta_1}) \times \\ &\left\{ P_1(t)(1 - t^{|m_1|}) + P_2(t)(1 - t^{\beta_1}) + P_3(t) \frac{1 - t^{|m_1|\beta_1}}{1 - t^{\beta_1}} + P_4(t) \frac{1 - t^{|m_1|\beta_1}}{1 - t^{|m_1|}} \right\} \\ &+ \sum_j G_j^1(t^{\beta_1})(1 - t^{|m_1|}) \times \left\{ P_5^j(t)(1 - t^{|m_1|}) + P_6^j(t)(1 - t^{\beta_1}) + \right. \\ &\left. P_7^j(t) \frac{1 - t^{|m_1|\beta_1}}{1 - t^{\beta_1}} + P_8^j(t) \frac{1 - t^{|m_1|\beta_1}}{1 - t^{|m_1|}} \right\} + (1 - t^{|m_1|})^{l_1-1} \times \\ &\left\{ P_9(t)(1 - t^{|m_1|}) + P_{10}(t)(1 - t^{\beta_1}) + P_{11}(t) \frac{1 - t^{|m_1|\beta_1}}{1 - t^{\beta_1}} + P_{12}(t) \frac{1 - t^{|m_1|\beta_1}}{1 - t^{|m_1|}} \right\}, \end{aligned}$$

where $P_k(t), P_k^j(t) \in \mathbb{Z}[t, t^{-1}]$ are Laurent polynomials. Since m_1 and β_1 are relatively prime, β_1 is odd. Substituting -1 for the above equation's t , we have

$$\Delta_{K_1}(-1)(2P_2(-1) + \beta_1 P_4(-1)) = \Delta_{K_2}((-1)^{\beta_2})(1 - (-1)^{|m_2|}).$$

If m_2 is even then $2P_2(-1) + \beta_1 P_4(-1) = 0$ since $\Delta_K(-1) \neq 0$ for any 1-knot K . If m_2 is odd then

$$\frac{\Delta_{K_2}(1)}{\Delta_{K_1}(-1)} = \frac{1}{\Delta_{K_1}(-1)} = \frac{2P_2(-1) + \beta_1 P_4(-1)}{2} \in \frac{\mathbb{Z}}{2}$$

since β_2 can be chosen to be even and $\Delta_K(1) = 1$ for any 1-knot K .

The same arguments for other generators $\Delta_{K_2}(t^{\beta_2}) \left\{ (1 - t^{\beta_2}), \frac{1 - t^{|m_2|/\beta_2}}{1 - t^{\beta_2}}, \frac{1 - t^{|m_2|/\beta_2}}{1 - t^{|m_2|}} \right\}$ of $E_1(K_2^{m_2, n_2})$ lead the following table:

The generators of $E_1(K_2^{m_2, n_2})$	$1 - t^{ m_2 }$	$1 - t^{\beta_2}$	$\frac{1 - t^{ m_2 /\beta_2}}{1 - t^{\beta_2}}$	$\frac{1 - t^{ m_2 /\beta_2}}{1 - t^{ m_2 }}$
$(m_2, \beta_1, \beta_2) = (e, o, o)$	P	$\mathbb{Z}/2$	P	\mathbb{Z}/β_2
$(m_2, \beta_1, \beta_2) = (o, o, e)$	$\mathbb{Z}/2$	P	$\mathbb{Z}/ m_2 $	P

The second column explains the case of the generator $\Delta_{K_2}(t^{\beta_2})(1 - t^{|m_2|})$, which we have seen above. If $(m_2, \beta_1, \beta_2) = (e, o, o)$, where e and o stands for even and odd, respectively, then we have $2P_2(-1) + \beta_1 P_4(-1) = 0$, which is represented ‘‘P’’ in the table. Note that we cannot get any information of $\Delta_{K_1}(t)$ and $\Delta_{K_2}(t)$ from the information ‘‘P’’. If $(m_2, \beta_1, \beta_2) = (o, o, e)$ then $\frac{\Delta_{K_2}((-1)^{\beta_2})}{\Delta_{K_1}(-1)} \in \mathbb{Z}/2$, which is represented by ‘‘ $\mathbb{Z}/2$ ’’. The 3rd, 4th, and 5th columns are filled by the same way for the other generators $\Delta_{K_2}(t^{\beta_2})(1 - t^{\beta_2})$, $\Delta_{K_2}(t^{\beta_2})\frac{1 - t^{|m_2|/\beta_2}}{1 - t^{\beta_2}}$, $\Delta_{K_2}(t^{\beta_2})\frac{1 - t^{|m_2|/\beta_2}}{1 - t^{|m_2|}}$, respectively.

In the case where m_2 is odd, from this table, we have $\frac{\Delta_{K_2}((-1)^{\beta_2})}{\Delta_{K_1}(-1)} \in \mathbb{Z}/2 \cap \mathbb{Z}/|m_2|$. We may choose β_2 to be even, then $|\Delta_{K_1}(-1)| = 1$ holds.

In the case where m_2 is even, since β_2 is odd, by the same argument we have $\frac{\Delta_{K_2}(-1)}{\Delta_{K_1}(-1)} \in \mathbb{Z}$. By applying the same argument with exchanging K_1 and K_2 , we have $\frac{\Delta_{K_1}(-1)}{\Delta_{K_2}(-1)} \in \mathbb{Z}$. Thus $|\Delta_{K_2}(-1)| = |\Delta_{K_1}(-1)|$. \square

REFERENCES

- [1] E. Artin, *Zur Isotopie zweidimensionalen Flächen im R^4* , Abh. Math. Sem. Univ. Hamburg **4** (1926), 47–72.
- [2] D. E. Blair, *Riemannian Geometry of Contact and Symplectic manifolds*, Progress in Mathematics, **203**, Birkhäuser Boston, Inc., Boston, 2002.
- [3] G. E. Bredon, *Introduction to compact transformation groups*, Academic Press, New York-London Pure and Applied Mathematics, Vol.46, 1972.
- [4] R. H. Cromwell and R. H. Fox, *Introduction to Knot Theory*, Graduate Texts in Math. No.57 Springer-Verlag, 1977.
- [5] R. Fintushel, *Locally smooth circle actions on homotopy 4-spheres*, Duke Math. J. **43** (1976), 63–70.
- [6] R. Jacoby, *One-parameter transformation groups of the three-sphere*, Proc. Amer. Math. Soc. **7** (1956), 131–140.
- [7] J. A. Hillman and S. P. Plotnick, *Geometrically fibered two-knots*, Math. Ann. **287** (1990) 259–273.

- [8] D. Montgomery and C. T. Yang, *Groups on S^n with principal orbits of dimension $n-3$* , **I, II**, Illinois J. Math. **4** (1960), 507–517. ; **5** (1961), 206–211.
- [9] P. S. Pao, *Non-linear circle actions on the 4-sphere and twisting spun knots*, Topology **17** (1978), 291–296.
- [10] S. Plotnick, *Fibered knots in S^4 -twisting, spinning, rolling, surgery, and branching*, Contemporary Math., vol. 5, Amer. Math. Soc., Providence, R.I., 1984, pp.437–459.
- [11] D. Rolfsen, *Knots and Links*, Math. Lec. Series, **7**, Publish or Perish, Inc., Berkeley, 1976.
- [12] M. Teragaito, *Roll-spun knots*, Math. Proc. Cambridge. Philos. Soc. **113**, no.1 (1992), 91–96.
- [13] M. Teragaito, *Twist-roll spun knots*, Proc, Amer, Math, Soc, **122**, no.2 (1994), 597-599.
- [14] E. C. Zeeman, *Twisting spun knots*, Trans. Am. math. Soc. **115** (1965), 471–495.

GRADUATE SCHOOL OF SCIENCE, TOHOKU UNIVERSITY, SENDAI, 980-8578, JAPAN

E-mail address: sb4m24@math.tohoku.ac.jp