

On five dimensional Sasakian Lie algebras with trivial center

E. Loiudice and A. Lotta

Abstract

We show that every five-dimensional Sasakian Lie algebra with trivial center is φ -symmetric. Moreover starting from a particular Sasakian structure on the Lie group $SL(2, \mathbb{R}) \times \text{Aff}(\mathbb{R})$ we obtain a family of contact metric (k, μ) structures whose Boeckx invariants assume all values less than -1 .

1 Introduction

In this paper we study some geometric features of Lie groups endowed with a left-invariant Sasakian structure. We focus on the five dimensional case, our basic tool being the complete classification of five dimensional Sasakian Lie algebras carried out by Andrada, Fino and Vezzoni in [1]. These Lie algebras fall into two classes: those whose center is one dimensional and generated by the Reeb vector field ξ , and those having trivial center.

Our results concern the centerless case, and our purpose is twofold. First of all we prove that, in this case, the simply connected Lie group G corresponding to a Sasakian Lie algebra is *always* a Sasakian φ -symmetric space. We remark that, in their classification of five dimensional Sasakian φ -symmetric Lie algebras carried out in [5], Calvaruso and Fino excluded the centerless Lie algebras, so our result fills this gap. Indeed, we study the canonical fibration $G \rightarrow G/H$ of the Sasakian space G over a homogeneous Kähler manifold G/H , showing that G/H is always either a product of two one dimensional complex space forms, namely $\mathbb{C}\mathbb{H}^1(\lambda) \times \mathbb{C}\mathbb{H}^1(\mu)$, or $\mathbb{C}\mathbb{P}^1(\lambda) \times \mathbb{C}\mathbb{H}^1(\mu)$, or it is a two dimensional complex space form $\mathbb{C}\mathbb{H}^2(\alpha)$.

Our second objective is to show that, starting from a particular Sasakian metric g on the Lie group $SL(2, \mathbb{R}) \times \text{Aff}(\mathbb{R})$, with contact form η , by a suitable deformation of g , one obtains a one parameter family of left-invariant metrics

Mathematics Subject Classification (2000): Primary 53C25, 22E60; Secondary 53C30, 53C35.

Keywords and Phrases: Sasakian Lie algebra, Sasakian φ -symmetric space, Contact metric (k, μ) -space.

g_a , $a > 1$, associated to the same η , for which the Reeb vector field ξ satisfies the (k, μ) curvature condition

$$R(X, Y)\xi = k(\eta(Y)X - \eta(X)Y) + \mu(\eta(Y)hX - \eta(X)hY), \quad k, \mu \in \mathbb{R}, \quad (1)$$

where $2h$ denotes the Lie derivative of the structural tensor φ along ξ . For more information about this widely studied class of associated metrics, we refer the reader to Blair's book [2, Chapter 7]. The Boeckx invariants I_a of these metrics exhaust all values in $(-\infty, -1)$. We observe that Boeckx himself in [3] provided Lie group models for contact metric (k, μ) spaces with $I < -1$; we feel that our construction provides an alternative way to obtain these models (at least in dimension five), which seems to be more geometrically transparent. Indeed, the definition of the metrics g_a relies on a general principle explored in [6], which provides (k, μ) structures starting from Sasakian ones endowed with additional geometric objects, namely a pair of conjugate, integrable Legendre distributions, whose Pang invariants are suitable multiples of the original Sasakian metric. However, no specific examples of application of these principle were given in [6], so we found interesting to carry out explicitly this construction on our Sasakian Lie groups.

2 Preliminaries

For general notions concerning contact metric geometry and for the notation we refer the reader to Blair's book [2]. It is well known that a locally symmetric Sasakian manifold must be a space of constant curvature [10]. This fact motivated Takahashi to introduce the notion of Sasakian φ -symmetric space, which is an analogous notion of Hermitian symmetric space, see [12]. We recall some basic facts concerning this concept. Let $(M, \varphi, \xi, \eta, g)$ be a Sasakian manifold. Consider an open neighborhood U of $x \in M$ such that the induced Sasakian structure on U , denoted with the same symbols, is regular. Let $\pi : U \rightarrow U/\xi$ be the corresponding fibration and (J, \bar{g}) the induced Kählerian structure on U/ξ [9]. Then M is called a *Sasakian locally φ -symmetric space* if U/ξ is always a Hermitian locally symmetric space.

This notion can be characterized also using the concept of φ -geodesic symmetry. A φ -geodesic of a Sasakian manifold $(M, \varphi, \xi, \eta, g)$ is a geodesic γ of M such that $\eta(\gamma') = 0$; while a φ -geodesic symmetry of M at $m \in M$ is a local diffeomorphism σ_m at m such that for each φ -geodesic $\gamma = \gamma(s)$, with $\gamma(0)$ in the trajectory of the integral curve of ξ passing through m , we have that

$$\sigma_m(\gamma(s)) = \gamma(-s)$$

for each s .

Theorem 1. *A Sasakian manifold M is a Sasakian locally φ -symmetric space if it admits at every point a φ -geodesic symmetry which is also a local automorphism of the structure.*

Let M be a Sasakian locally φ -symmetric space. If any φ -geodesic symmetry of M is extendable to a global automorphism of M , and if the Killing vector field ξ generates a one parameter group of global transformations, then M is called a *Sasakian φ -symmetric space*. The following theorems are proved in [12].

Theorem 2. *A Sasakian globally φ -symmetric space is a principal G^1 -bundle over a Hermitian globally symmetric space.*

Here G^1 is a 1-dimensional Lie group isomorphic to the one parameter group of transformations generated by ξ .

Theorem 3. *A complete and simply connected Sasakian locally φ -symmetric space is a globally φ -symmetric space.*

Now let G be a simply connected Lie group endowed with a left-invariant Sasakian structure (φ, ξ, η, g) . Let \mathfrak{g} be the Lie algebra of G , so that $\xi \in \mathfrak{g}$. Then, by homogeneity, ξ is regular (cf. [4, Theorem 4]) and the fibration of G is the canonical projection $p : G \rightarrow G/H$, where $H \subset G$ is the analytic subgroup with $\text{Lie}(H) = \mathbb{R}\xi$. In this case, the induced Kähler metric \bar{g} on G/H is the unique one making p a Riemannian submersion; clearly it is G -invariant. Hence G is a Sasakian φ -symmetric space if and only if G/H is a Hermitian symmetric space.

3 Five-dimensional Sasakian Lie algebras

A *Sasakian Lie algebra* is a Lie algebra \mathfrak{g} endowed with a quadruple (φ, ξ, η, g) where $\varphi \in \text{End}(\mathfrak{g})$, $\xi \in \mathfrak{g}$, $\eta \in \mathfrak{g}^*$ and g is an inner product such that

$$\begin{aligned} \eta(\xi) = 1, \quad \varphi^2 = -Id + \eta \otimes \xi, \quad g(\varphi X, \varphi Y) = g(X, Y) - \eta(X)\eta(Y), \\ g(X, \varphi Y) = d\eta(X, Y), \quad N_\varphi = -2d\eta \otimes \xi, \end{aligned} \quad (2)$$

where

$$N_\varphi(X, Y) := \varphi^2[X, Y] + [\varphi X, \varphi Y] - \varphi[\varphi X, Y] - \varphi[X, \varphi Y].$$

Sasakian Lie algebras have been classified in [1] treating separately the two cases of Lie algebras with trivial or nontrivial center. We recall the classification result concerning Sasakian Lie algebras with trivial center [1, Theorem 13]:

Theorem 4. *Let $(\mathfrak{g}, \varphi, \xi, \eta, g)$ be a five dimensional Sasakian Lie algebra with trivial center. Then \mathfrak{g} is isomorphic to one of the following Lie algebras: the direct products $\mathfrak{sl}(2, \mathbb{R}) \times \mathfrak{aff}(\mathbb{R})$, $\mathfrak{su}(2) \times \mathfrak{aff}(\mathbb{R})$, or the non-unimodular solvable Lie algebra $\mathbb{R}^2 \ltimes \mathfrak{h}_3$.*

Here $\mathfrak{aff}(\mathbb{R})$ and \mathfrak{h}_3 denote respectively the Lie algebra of the Lie group of affine motions of \mathbb{R} and the real three-dimensional Heisenberg Lie algebra.

Let $(\mathfrak{g}, \varphi, \xi, \eta, g)$ be a five dimensional Sasakian Lie algebra with trivial center. According to [1, Section 3.2], after performing a \mathcal{D} -homothetic deformation of the structure, there exists an orthonormal basis $\{e_1, \dots, e_5\}$ of \mathfrak{g} satisfying:

$$\ker \eta = \text{span}\{e_1, \dots, e_4\}, \quad e_5 = \xi, \quad \varphi(e_1) = -e_2, \quad \varphi(e_3) = -e_4,$$

and such that one of the following eight sets of bracket relations holds:

$$\begin{aligned} [e_1, e_2] &= \frac{2}{c}e_1 - 2e_5, & [e_1, e_3] &= ce_4, & [e_1, e_4] &= -ce_3, & [e_1, e_5] &= 0, \\ [e_2, e_3] &= -fe_4, & [e_2, e_4] &= fe_3, & [e_2, e_5] &= 0, & [e_3, e_4] &= -2e_5, \\ [e_3, e_5] &= -e_4, & [e_4, e_5] &= e_3, \end{aligned} \quad (\text{A1})$$

$$\begin{aligned} [e_1, e_2] &= -ae_1 - be_2 - 2e_5, & [e_1, e_3] &= ce_4, & [e_1, e_4] &= -ce_3, \\ [e_1, e_5] &= 0, & [e_2, e_3] &= -\frac{2+ac}{b}e_4, & [e_2, e_4] &= \frac{2+ac}{b}e_3, \\ [e_2, e_5] &= 0, & [e_3, e_4] &= -2e_5, & [e_3, e_5] &= -e_4, \\ [e_4, e_5] &= e_3, \end{aligned} \quad (\text{A2})$$

$$\begin{aligned} [e_1, e_2] &= -ae_1 - 2e_5, & [e_1, e_3] &= \frac{2}{a}e_4, & [e_1, e_4] &= -\frac{2}{a}e_3, & [e_1, e_5] &= 0, \\ [e_2, e_3] &= -fe_4, & [e_2, e_4] &= fe_3, & [e_3, e_4] &= -2e_5, & [e_2, e_5] &= 0, \\ [e_3, e_5] &= e_4, & [e_4, e_5] &= -e_3, \end{aligned} \quad (\text{B1})$$

$$\begin{aligned} [e_1, e_2] &= -ae_1 - be_2 - 2e_5, & [e_1, e_3] &= ce_4, & [e_1, e_4] &= -ce_3, \\ [e_1, e_5] &= 0, & [e_2, e_3] &= -\frac{ac-2}{b}e_4, & [e_2, e_4] &= \frac{ac-2}{b}e_3, \\ [e_2, e_5] &= 0, & [e_3, e_4] &= -2e_5, & [e_3, e_5] &= e_4, \\ [e_4, e_5] &= -e_3, \end{aligned} \quad (\text{B2})$$

$$\begin{aligned} [e_1, e_2] &= -ae_1 - 2e_5, & [e_1, e_3] &= -\frac{2}{a}e_4, & [e_1, e_4] &= \frac{2}{a}e_3, \\ [e_1, e_5] &= 0, & [e_2, e_3] &= \frac{a}{2}e_3 - fe_4, & [e_2, e_4] &= fe_3 + \frac{a}{2}e_4, \\ [e_2, e_5] &= 0, & [e_3, e_4] &= -ae_1 - 2e_5, & [e_3, e_5] &= -e_4, \\ [e_4, e_5] &= e_3, \end{aligned} \quad (\text{A3})$$

$$\begin{aligned} [e_1, e_2] &= -ae_1 - be_2 - 2e_5, & [e_1, e_3] &= -\frac{1}{2}be_3 + ce_4, \\ [e_1, e_4] &= -ce_3 - \frac{1}{2}be_4, & [e_1, e_5] &= 0, \\ [e_2, e_3] &= \frac{1}{2}ae_3 - \frac{2+ac}{b}e_4, & [e_2, e_4] &= \frac{2+ac}{b}e_3 + \frac{1}{2}ae_4, \\ [e_2, e_5] &= 0, & [e_3, e_4] &= -ae_1 - be_2 - 2e_5, \\ [e_3, e_5] &= -e_4, & [e_4, e_5] &= e_3, \end{aligned} \quad (\text{A4})$$

$$\begin{aligned}
[e_1, e_2] &= -ae_1 - 2e_5, & [e_1, e_3] &= \frac{2}{a}e_4, & [e_1, e_4] &= -\frac{2}{a}e_3, \\
[e_1, e_5] &= 0, & [e_2, e_3] &= \frac{a}{2}e_3 - fe_4, & [e_2, e_4] &= fe_3 + \frac{a}{2}e_4, \\
[e_2, e_5] &= 0, & [e_3, e_4] &= -ae_1 - 2e_5, & [e_3, e_5] &= e_4, \\
[e_4, e_5] &= -e_3,
\end{aligned} \tag{B3}$$

$$\begin{aligned}
[e_1, e_2] &= -ae_1 - be_2 - 2e_5, & [e_1, e_3] &= -\frac{1}{2}be_3 + ce_4, \\
[e_1, e_4] &= -ce_3 - \frac{1}{2}be_4, & [e_1, e_5] &= 0, \\
[e_2, e_3] &= \frac{1}{2}ae_3 - \frac{ac-2}{b}e_4, & [e_2, e_4] &= \frac{ac-2}{b}e_3 + \frac{1}{2}ae_4, \\
[e_2, e_5] &= 0, & [e_3, e_4] &= -ae_1 - be_2 - 2e_5, \\
[e_4, e_5] &= -e_3, & [e_3, e_5] &= e_4.
\end{aligned} \tag{B4}$$

In the first two cases we have that the Lie algebra \mathfrak{g} is isomorphic to the direct product $\mathfrak{sl}(2, \mathbb{R}) \times \mathfrak{aff}(\mathbb{R})$, where

$$\begin{cases} \mathfrak{aff}(\mathbb{R}) \simeq \text{span}\{fe_1 + ce_2, e_1 + ce_5\} \\ \mathfrak{sl}(2, \mathbb{R}) \simeq \text{span}\{e_3, e_4, e_5\} \end{cases}$$

in case (A1), and

$$\begin{cases} \mathfrak{aff}(\mathbb{R}) \simeq \text{span}\{ae_1 + be_2 + 2e_5, e_1 - ce_5\} \\ \mathfrak{sl}(2, \mathbb{R}) \simeq \text{span}\{e_3, e_4, e_5\} \end{cases}$$

in case (A2).

In cases (B1) and (B2) we have respectively that

$$\begin{cases} \mathfrak{aff}(\mathbb{R}) \simeq \text{span}\{ae_1 + e_5, fe_1 + e_5\} \\ \mathfrak{su}(2) \simeq \text{span}\{e_3, e_4, e_5\} \end{cases}$$

or

$$\begin{cases} \mathfrak{aff}(\mathbb{R}) \simeq \text{span}\{ae_1 + be_2 + 2e_5, e_1 + ce_5\} \\ \mathfrak{su}(2) \simeq \text{span}\{e_3, e_4, e_5\} \end{cases}$$

and thus \mathfrak{g} is isomorphic to the direct product $\mathfrak{su}(2) \times \mathfrak{aff}(\mathbb{R})$. Lastly, in cases (A3), (A4), (B3), (B4) one has $\mathfrak{g} \simeq \mathbb{R}^2 \times \mathfrak{h}_3$.

4 Sasakian φ -symmetric five-dimensional Lie groups

In this section we prove our main result:

Theorem 5. *Let G be a five-dimensional simply connected Lie group endowed with a left-invariant Sasakian structure. If the Lie algebra of G has trivial center, then G is a Sasakian φ -symmetric space.*

We need the following:

Lemma 1. *Let $(G/H, g)$ be a homogeneous Riemannian manifold, where G is a connected Lie group and suppose we are given a reductive decomposition:*

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m},$$

of the Lie algebra \mathfrak{g} of G . We suppose there exists an $Ad(H)$ -invariant subspace \mathfrak{n} of \mathfrak{m} satisfying:

$$[\mathfrak{g}, \mathfrak{h} \oplus \mathfrak{n}] \subset \mathfrak{h} \oplus \mathfrak{n}, \quad U(\mathfrak{m}, \mathfrak{n}) \subset \mathfrak{n}, \quad (3)$$

where $U : \mathfrak{m} \times \mathfrak{m} \rightarrow \mathfrak{m}$ is the bilinear operator defined by

$$2\langle U(X, Y), Z \rangle = \langle [Z, X]_{\mathfrak{m}}, Y \rangle + \langle [Z, Y]_{\mathfrak{m}}, X \rangle.$$

Here $\langle \cdot, \cdot \rangle$ is the $Ad(H)$ -invariant scalar product on \mathfrak{m} corresponding to the Riemannian metric g on G/H . Then the G -invariant distribution \mathcal{D} determined by \mathfrak{n} is parallel with respect to the Levi-Civita connection of g .

Proof. Let $\pi : G \rightarrow G/H$ be the canonical projection, $e \in G$ the identity element of G and $\varrho = \pi(e)$. For every $a \in G$ we denote by τ_a the isometry of G/H given by $\tau_a(bH) = abH$. First of all we show that the fundamental vector field X^* determined by a vector $X \in \mathfrak{n}$ is a section of \mathcal{D} . Indeed, for every $a \in G$, according to (3) we have:

$$(Ad(a^{-1})X) \in \mathfrak{h} \oplus \mathfrak{n},$$

hence

$$(d\tau_{a^{-1}})_{aH}(X^*_{aH}) = (Ad(a^{-1})X)_{\varrho}^* = (d\pi)_e(Ad(a^{-1})X) \in \mathcal{D}_{\varrho},$$

and this ensures that $X^*_{aH} \in \mathcal{D}_{aH}$. Now, let $\{e_1, \dots, e_k\}$ be a basis of \mathfrak{n} . Then

$$\{(e_1^*)_{aH}, \dots, (e_k^*)_{aH}\}$$

is a basis of \mathcal{D}_{aH} , for every point aH in a neighborhood W of ϱ . Using the standard formula for the Levi-Civita connection of a homogeneous Riemannian reductive space [8], and the assumption (3), we get:

$$\nabla_X e_i^* = -\frac{1}{2}[X, e_i]_{\mathfrak{m}} + U(X, e_i) \in \mathfrak{n},$$

for every $X \in T_{\varrho}(G/H)$. Here we are identifying $T_{\varrho}(G/H)$ with \mathfrak{m} via $(d\pi)_e|_{\mathfrak{m}} : \mathfrak{m} \rightarrow T_{\varrho}(G/H)$. Then we see that for every section Y of \mathcal{D} and $X \in T_{\varrho}(G/H)$, $\nabla_X Y$ belongs to \mathcal{D}_{ϱ} . Hence, by the G -invariance, we can conclude that \mathcal{D} is parallel. \square

Proof of the theorem. Let (φ, ξ, η, g) be a left-invariant Sasakian structure on G . We denote by H the one parameter subgroup generated by ξ , and with \bar{g} the induced Kähler metric on G/H via the canonical submersion $\pi : G \rightarrow G/H$. We denote by $\mathfrak{h} = \mathbb{R}\xi$ the Lie algebra of H . As we recalled in Section 3, up to a \mathcal{D} -homothetic deformation, there exists an orthonormal basis $\{e_1, \dots, e_5\}$ of the Lie algebra \mathfrak{g} of G such that

$$\ker \eta = \text{span}\{e_1, \dots, e_4\}, \quad e_5 = \xi, \quad \varphi(e_1) = -e_2, \quad \varphi(e_3) = -e_4,$$

and the elements e_1, \dots, e_5 satisfy one of the bracket relations (A1)-(A4) or (B1)-(B4). Moreover, G/H is a homogeneous reductive space with a reductive decomposition given by

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m},$$

where $\mathfrak{m} := \text{span}\{e_1, \dots, e_4\}$. The Kähler metric \bar{g} on G/H is the G -invariant metric induced by the scalar product g_e on \mathfrak{m} .

In cases (A1), (A2), (B1), (B2) we consider the G -invariant distributions \mathcal{D}_1 and \mathcal{D}_2 on G/H determined respectively by

$$\mathfrak{n}_1 := \text{span}\{e_1, e_2\}, \quad \mathfrak{n}_2 := \text{span}\{e_3, e_4\}.$$

Then \mathcal{D}_1 and \mathcal{D}_2 are orthogonal and complementary distributions on G/H and one can easily check directly that \mathfrak{n}_2 satisfies the conditions (3) of Lemma 1, yielding that \mathcal{D}_2 is parallel. For example, in case (A1) we have:

$$\begin{aligned} U(e_3, e_1) &= \frac{1}{2}ce_4, & U(e_3, e_2) &= -\frac{1}{2}fe_4, & U(e_3, e_3) &= 0, & U(e_3, e_4) &= 0, \\ U(e_4, e_1) &= -\frac{1}{2}ce_3, & U(e_4, e_2) &= \frac{1}{2}fe_3, & U(e_4, e_4) &= 0. \end{aligned}$$

Then G/H is isometric to the direct product $M_1 \times M_2$, where M_1 and M_2 are both 2-dimensional homogeneous Kähler manifolds (cf. [8, Theorem 5.1, p. 211 and Theorem 8.1, p. 172]). Of course, each of these spaces has constant curvature, and this proves that G/H is a Hermitian symmetric space. Actually in case (A1) and (A2) the sectional curvature of M_1 is $\lambda = -\frac{4}{c^2}$ or $\lambda = -(a^2 + b^2)$ respectively, and the sectional curvature of M_2 is $\mu = -2$ in both cases; while $\lambda = -a^2$, $\mu = 2$ in case (B1) and $\lambda = -(a^2 + b^2)$, $\mu = 2$ in case (B2), as can be readily checked explicitly by using the corresponding bracket relations. Of course, this also yields that, as an Hermitian symmetric space, we have that

$$G/H \cong \mathbb{C}\mathbb{H}^1(\lambda) \times \mathbb{C}\mathbb{H}^1(\mu)$$

in cases (A1) and (A2), or

$$G/H \cong \mathbb{C}\mathbb{H}^1(\lambda) \times \mathbb{C}\mathbb{P}^1(\mu)$$

in cases (B1) and (B2).

Now consider the case (A3). We set:

$$E_1 = ae_1 + 2e_5, \quad E_2 = \frac{1}{a}e_2, \quad E_3 = e_3, \quad E_4 = e_4,$$

and $\mathfrak{s} := \text{span}\{E_1, \dots, E_4\}$. Then \mathfrak{s} is an ideal of \mathfrak{g} and

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{s}.$$

Hence the analytic subgroup S of G corresponding to \mathfrak{s} acts simply transitively on G/H and the restriction $p : S \rightarrow G/H$ provides a diffeomorphism between S and G/H . The G -invariant Kähler structure (J, \bar{g}) of G/H transfers via p to a left-invariant Kähler structure on S , denoted by the same symbols. We claim that S is a complex hyperbolic space $\mathbb{C}H^2(\alpha)$.

Observe that J is determined on \mathfrak{s} by:

$$JE_1 = -a^2E_2, \quad JE_2 = \frac{1}{a^2}E_1, \quad JE_3 = -E_4, \quad JE_4 = E_3.$$

Moreover, \mathfrak{s} admits the following \bar{g} -orthogonal decomposition:

$$\mathfrak{s} = \mathbb{R}A_0 \oplus \mathfrak{a}_1 \oplus \mathfrak{a}_2,$$

where

$$A_0 = |a|E_2, \quad \mathfrak{a}_1 = \text{span}\{E_3, E_4\}, \quad \mathfrak{a}_2 = \text{span}\{JA_0\}, \quad \lambda = \frac{|a|}{2}.$$

The following relations hold:

$$\begin{aligned} [A_0, X] &= \frac{|a|}{2}X + S_0(X), & [A_0, JA_0] &= |a|JA_0 \\ [X, Y] &= |a|\bar{g}(JX, Y)JA_0, & [X, JA_0] &= 0, \end{aligned}$$

for every $X, Y \in \mathfrak{a}_1$, where $S_0 : \mathfrak{a}_1 + \mathfrak{a}_2 \rightarrow \mathfrak{a}_1 + \mathfrak{a}_2$ is the skew-symmetric derivation defined by

$$S_0(E_3) = -\frac{af}{|a|}E_4, \quad S_0(E_4) = \frac{af}{|a|}E_3, \quad S_0(JA_0) = 0.$$

Now our claim follows from Heintze's description of the complex hyperbolic space as a solvable Lie group endowed with a left-invariant Kähler metric of negative curvature (cf. [7, Section 6]).

All the remaining cases can be treated similarly. In case (A4) consider

$$E_1 = ae_1 + be_2 + 2e_5, \quad E_2 = \frac{1}{b}e_1, \quad E_3 = e_3, \quad E_4 = e_4,$$

and again the ideal $\mathfrak{s} = \text{span}\{E_1, \dots, E_4\}$. Now we have:

$$JE_1 = -\frac{a}{b}E_1 + (a^2 + b^2)E_2, \quad JE_2 = -\frac{1}{b^2}E_1 + \frac{a}{b}E_2, \quad JE_3 = -E_4, \quad JE_4 = E_3,$$

and \mathfrak{s} admits the orthogonal decomposition:

$$\mathfrak{s} = \mathbb{R}A_0 \oplus \mathfrak{a}_1 \oplus \mathfrak{a}_2,$$

with

$$\begin{aligned} A_0 &= \frac{1}{2\lambda} \left(\frac{a}{b} E_1 - (a^2 + b^2) E_2 \right), \quad \lambda = \sqrt{\frac{a^2 + b^2}{4}} \\ \mathfrak{a}_1 &= \text{span}\{E_3, E_4\}, \quad \mathfrak{a}_2 = \text{span}\{JA_0\}, \end{aligned} \quad (4)$$

Then the following relations hold

$$\begin{aligned} [A_0, X] &= \lambda X + S_0(X), & [A_0, JA_0] &= 2\lambda JA_0 \\ [X, Y] &= 2\lambda \bar{g}(JX, Y) JA_0, & [X, JA_0] &= 0, \end{aligned}$$

for $X, Y \in \mathfrak{a}_1$, where $S_0 : \mathfrak{a}_1 + \mathfrak{a}_2 \rightarrow \mathfrak{a}_1 + \mathfrak{a}_2$ is now given by

$$S_0(E_3) = -2\lambda \frac{c}{b} E_4, \quad S_0(E_4) = 2\lambda \frac{c}{b} E_3, \quad S_0(JA_0) = 0,$$

and thus again S is a complex hyperbolic space. Cases (B3) and (B4) are referable to cases (A3) and (A4) respectively. Indeed we can consider the ideals

$$\mathfrak{s} = \text{span}\left\{ae_1 + 2e_5, \frac{1}{a}e_2, e_3, e_4\right\}$$

in case (B3) and

$$\mathfrak{s} = \text{span}\left\{ae_1 + be_2 + 2e_5, \frac{1}{b}e_1, e_3, e_4\right\}$$

in case (B4), and the left-invariant Kähler structure on such Lie algebras obtained as before. One can check that in the first case this Lie algebra \mathfrak{s} is isometric to the one considered in case (A3), while in the second case \mathfrak{s} is isometric to the Lie algebra considered in case (A4), thus completing the proof. \square

5 (k, μ) -contact metric spaces of dimension five

In [3] Boeckx classified all the non-Sasakian contact metric (k, μ) -spaces, i.e., the contact metric manifolds whose curvature satisfies condition (1), up to \mathcal{D} -homothetic transformations. He showed that such a space is locally determined by its dimension and the number $I = \frac{1-\mu}{\sqrt{1-k}}$; moreover he gave explicit examples of (k, μ) -spaces, consisting of some abstract Lie groups, for every possible dimension and for every $I \leq -1$. We remark that in his exposition Boeckx does not give any explicit description of these Lie groups. In this section we shall construct 5-dimensional non-Sasakian (k, μ) -spaces with Boeckx invariant $I < -1$ in terms of classical Lie groups, using the classification of the 5-dimensional Sasakian Lie algebras described in section 3. Our tool will be the relation between Sasakian structures and non-Sasakian (k, μ) -structures

explored by Cappelletti-Montano, Carriazo and Martín-Molina in [6]. Their approach is based on the Pang invariant of a Legendre foliation (cf. [11]). We recall that, if \mathcal{D} is an integrable Legendre distribution on a contact manifold (M, η) , its Pang invariant is the symmetric bilinear form on the tangent bundle of \mathcal{D} , defined by

$$\Pi_{\mathcal{D}}(X, X') = -(\mathcal{L}_X \mathcal{L}_{X'} \eta)(\xi) = 2d\eta([\xi, X], X'),$$

where ξ is the Reeb vector field.

Theorem 6. *Let \mathfrak{g} be the Sasakian Lie algebra of type (A2) with*

$$c = a = 0, \quad b = -\sqrt{2}.$$

Let G be a connected Lie group with Lie algebra \mathfrak{g} and denote by (φ, ξ, η, g) the associated left-invariant Sasakian structure on G . Then \mathfrak{g} admits an orthogonal decomposition

$$\mathfrak{g} = \mathfrak{d}_+ \oplus \mathfrak{d}_- \oplus \mathbb{R}\xi$$

such that, for every real number $a > 1$, the modified left-invariant metric g_a defined by:

$$(g_a)_e = \begin{cases} \frac{1}{a}g_e, & \text{on } \mathfrak{d}_+ \times \mathfrak{d}_+ \\ ag_e, & \text{on } \mathfrak{d}_- \times \mathfrak{d}_-, \\ \eta_e \otimes \eta_e, & \text{otherwise} \end{cases}$$

is also associated to the contact form η and determines a contact metric (k, μ) structure on G , whose Boeckx invariant is

$$I_a = -\frac{a^2 + 1}{a^2 - 1}. \quad (5)$$

Proof. Let $\{e_1, \dots, e_5\}$ be the orthonormal basis of the Sasakian Lie algebra $(\mathfrak{g}, \varphi, \xi, \eta, g)$, with

$$\ker \eta = \text{span}\{e_1, \dots, e_4\}, \quad e_5 = \xi, \quad \varphi(e_1) = -e_2, \quad \varphi(e_3) = -e_4, \quad (6)$$

and for which the nonzero bracket relations are:

$$\begin{aligned} [e_1, e_2] &= \sqrt{2}e_2 - 2e_5, & [e_2, e_3] &= \sqrt{2}e_4, & [e_2, e_4] &= -\sqrt{2}e_3, \\ [e_4, e_5] &= e_3, & [e_3, e_4] &= -2e_5, & [e_3, e_5] &= -e_4. \end{aligned}$$

Now, we observe that the following two Lie sub-algebras of \mathfrak{g} :

$$\mathfrak{d}_+ := \text{span}\{e_1 + e_4, e_2 + e_3\}, \quad \mathfrak{d}_- := \text{span}\{e_3 - e_2, e_1 - e_4\}$$

determine two totally geodesic, mutually orthogonal, Legendre left-invariant foliations \mathcal{D}_+ and \mathcal{D}_- of G with respect to the Sasakian structure. Moreover, the Pang invariants $\Pi_{\mathcal{D}_+}$ and $\Pi_{\mathcal{D}_-}$ are determined at the neutral element of G by:

$$\begin{aligned} \Pi_{\mathcal{D}_+}(X, Y) &= 2d\eta([\xi, X], Y) = -g(X, Y), \\ \Pi_{\mathcal{D}_-}(V, W) &= 2d\eta([\xi, V], W) = -g(V, W), \end{aligned}$$

where $X, Y \in \mathfrak{d}_+$ and $V, W \in \mathfrak{d}_-$. Hence, by the left-invariance of the Sasakian structure, Theorem 4.1 of [6] is applicable, ensuring that there exists a family of contact metrics (k_a, μ_a) -structure on G compatible with the contact form η , with

$$g_a = \begin{cases} -\frac{1}{a}g, & \text{on } T\mathcal{D}_+ \times T\mathcal{D}_+ \\ -ag, & \text{on } T\mathcal{D}_- \times T\mathcal{D}_- \\ \eta \otimes \eta, & \text{otherwise} \end{cases}, \quad \varphi_a = \begin{cases} -\frac{1}{a}\varphi, & \text{on } T\mathcal{D}_+ \\ -a\varphi, & \text{on } T\mathcal{D}_- \\ 0, & \text{on } \mathbb{R}\xi \end{cases}, \quad (7)$$

$$k_a = 1 - \frac{(a^2 - 1)^2}{16a^2}, \quad \mu_a = 2 - \frac{a^2 + 1}{2a},$$

and $I_a = -\frac{a^2+1}{|a^2-1|}$, parametrized by $a < -1$. This yields the result. \square

Since I_a in (5) assumes all values in $(-\infty, -1)$, we can reformulate the above result in order to obtain the following local classification of five dimensional (k, μ) contact metric spaces. Denote by $\{E_1, \dots, E_5\}$ the standard basis of the Lie algebra $\mathfrak{sl}(2, \mathbb{R}) \times \mathfrak{aff}(\mathbb{R})$, satisfying the following relations:

$$[E_1, E_2] = E_2, \quad [E_3, E_4] = -E_5, \quad [E_3, E_5] = -2E_4, \quad [E_4, E_5] = 2E_3.$$

Corollary 1. *Every five-dimensional contact metric (k, μ) -space with Boeckx invariant $I < -1$ is locally equivalent, up to a \mathcal{D} -homothetic deformation, to the direct product*

$$SL(2, \mathbb{R}) \times Aff(\mathbb{R}),$$

endowed with the left invariant contact metric structure (φ, ξ, η, g) determined with respect to the standard basis $\{E_1, \dots, E_5\}$ of $\mathfrak{aff}(\mathbb{R}) \times \mathfrak{sl}(2, \mathbb{R})$ by:

$$\varphi \equiv \begin{pmatrix} 0 & 2s & t & 0 & 0 \\ -\frac{1}{2}s & 0 & 0 & \frac{1}{2}t & 0 \\ \frac{1}{2}t & 0 & 0 & -s & 0 \\ 0 & t & s & 0 & 0 \\ \frac{1}{2}s & 0 & 0 & -\frac{1}{2}t & 0 \end{pmatrix}, \quad g \equiv \begin{pmatrix} -\frac{1}{2}s & 0 & 0 & \frac{1}{2}t & 0 \\ 0 & 4 - 2s & -t & 0 & 4 \\ 0 & -t & -s & 0 & 0 \\ \frac{1}{2}t & 0 & 0 & -s & 0 \\ 0 & 4 & 0 & 0 & 4 \end{pmatrix},$$

$$\eta(E_1) = \eta(E_3) = \eta(E_4) = 0, \quad \eta(E_2) = \eta(E_5) = 2, \quad \xi = \frac{1}{2}E_5,$$

where

$$s = -\frac{1}{2}\left(\sqrt{\frac{I-1}{I+1}} + \sqrt{\frac{I+1}{I-1}}\right), \quad t = -\frac{1}{\sqrt{2}}\left(\sqrt{\frac{I-1}{I+1}} - \sqrt{\frac{I+1}{I-1}}\right).$$

Remark 1. Concerning the already cited examples of (k, μ) -spaces provided by Boeckx, they consist in a 2-parameter family of abstract Lie groups, endowed with a left-invariant contact metric structure. Actually, one can check that, in the five dimensional case, the corresponding Lie algebras are all isomorphic to $\mathfrak{sl}(2, \mathbb{R}) \times \mathfrak{aff}(\mathbb{R})$.

References

- [1] A. Andrada, A. Fino, and L. Vezzoni: *A class of Sasakian 5-manifolds*, Transform. Groups **14** (2009), 493–512.
- [2] D. E. Blair: *Riemannian geometry of contact and symplectic manifolds*, volume 203 of *Progress in Mathematics*. Birkhäuser, Boston Inc., Boston, MA, second edition, 2010.
- [3] E. Boeckx: *A full classification of contact metric (k, μ) -spaces*, Illinois J. Math. **44** (2000), 212–219.
- [4] W. M. Boothby and H. C. Wang: *On contact manifolds*, Ann. of Math. (2) **68** (1958), 721–734.
- [5] G. Calvaruso and A. Fino: *Five-dimensional K -contact Lie algebras*, Monatsh. Math. **167** (2012), 35–59.
- [6] B. Cappelletti-Montano, A. Carriazo, and V. Martín-Molina: *Sasaki-Einstein and paraSasaki-Einstein metrics from (k, μ) -structures*, J. Geom. Phys. **73** (2013), 20–36.
- [7] E. Heintze: *On homogeneous manifolds of negative curvature*, Math. Ann. **211** (1974), 23–34.
- [8] S. Kobayashi and K Nomizu: *Foundations of differential geometry. Vol. II*, Interscience Publishers, 1969.
- [9] K. Ogiue: *On fiberings of almost contact manifolds*, Kōdai Math. Sem. Rep. **17** (1965), 53–62.
- [10] M. Okumura: *Some remarks on space with a certain contact structure*, Tōhoku Math. J. (2) **14** (1962), 135–145.
- [11] M. Y. Pang: *The structure of Legendre foliations*, Trans. Amer. Math. Soc. **320** (1990), 417–455.
- [12] T. Takahashi: *Sasakian ϕ -symmetric spaces*, Tōhoku Math. J. (2) **29** (1977), 91–113.

Eugenia Loiudice

Dipartimento di Matematica, Università di Bari Aldo Moro,
Via Orabona 4, 70125 Bari, Italy
e-mail: eugenia.loiudice@uniba.it

Antonio Lotta

Dipartimento di Matematica, Università di Bari Aldo Moro,
Via Orabona 4, 70125 Bari, Italy
e-mail: antonio.lotta@uniba.it